HELCOM Monitoring Programme topic

Seabed habitat distribution and extent

Programme:

Habitat-forming species and substrates

Contents

a.	Metadata on monitoring strategies and monitoring programmes	2
	a.1 Responsible HELCOM subsidiary body	2
	a.2 Regional Cooperation (RegionalCooperation)	3
b.	Monitoring strategies	3
	b.1 Descriptor	3
	b.2 BSAP segments	4
	b.3 Monitoring strategy description	4
	b.4 BSAP Ecological objectives	4
	b.5 Gaps in monitoring	5
c.	Monitoring programmes	5
	c.1 Purpose of monitoring	5
	c.2 Other legislation	9
	c.3 Implementation of Regional Cooperation (RegionalCooperation_implementation)	. 10
	c.4 Monitoring concepts	. 11
	c.5 Monitoring and assessment requirements	. 16
	c.6 Data providers and access	. 17
	c.7 MSFD Criteria (GES Criteria)	. 20
٦	References	27

a. Metadata on monitoring strategies and monitoring programmes

a.1 Responsible HELCOM subsidiary body

Please indicate the relevant expert group/network if available, otherwise the responsible HELCOM Working Group.

	Permament Groups
	Gear – Group on the Implementation of the Ecosystem Approach
	Maritime – Maritime Working Group
	Pressure – Working Group on Reduction of Pressures from the Baltic Sea Catchment Area
	Response – Response Working Group
\boxtimes	State and Conservation – Working Group on the State of the Environmental and Natgure Conservation
	Time-limited Groups
	Agri – Group on Sustainable Agricultural Practices
	Fish – Group on Ecosystem-based Sustainable Fisheries
	HELCOM-VASAB MSP WG - Joint HELCOM-VASAB Maritime Spatial Planning Working Group
	Expert Groups
	Expert Groups AIS EWG – Expert Working Group for Mutual Exchange and Deliveries of AIS data
	AIS EWG – Expert Working Group for Mutual Exchange and Deliveries of AIS data
	AIS EWG – Expert Working Group for Mutual Exchange and Deliveries of AIS data EN Hazardous Substances – Expert Network on hazardous substances
	AIS EWG – Expert Working Group for Mutual Exchange and Deliveries of AIS data EN Hazardous Substances – Expert Network on hazardous substances EN Marine Litter – Expert Network on Marine Litter
	AIS EWG – Expert Working Group for Mutual Exchange and Deliveries of AIS data EN Hazardous Substances – Expert Network on hazardous substances EN Marine Litter – Expert Network on Marine Litter EN Noise – Expert Network on Underwater Noise
	AIS EWG – Expert Working Group for Mutual Exchange and Deliveries of AIS data EN Hazardous Substances – Expert Network on hazardous substances EN Marine Litter – Expert Network on Marine Litter EN Noise – Expert Network on Underwater Noise ESA – Expert Network on Economic and Social Analyses
	AIS EWG – Expert Working Group for Mutual Exchange and Deliveries of AIS data EN Hazardous Substances – Expert Network on hazardous substances EN Marine Litter – Expert Network on Marine Litter EN Noise – Expert Network on Underwater Noise ESA – Expert Network on Economic and Social Analyses EWG OWR – Expert Working Group on Oiled Wildlife Response
	AIS EWG – Expert Working Group for Mutual Exchange and Deliveries of AIS data EN Hazardous Substances – Expert Network on hazardous substances EN Marine Litter – Expert Network on Marine Litter EN Noise – Expert Network on Underwater Noise ESA – Expert Network on Economic and Social Analyses EWG OWR – Expert Working Group on Oiled Wildlife Response EWG SHORE – Expert Working Group on Response on the Shore

	IN-EUTROPHICATION - Intersessional Network on Eutrophication
	IWGAS – Informal Working Group on Aerial Surveillance
	JWG Bird – HELCOM-OSPAR-ICES Joint Working Group on Seabirds
	MORS EG – Expert group on monitoring of radioactive substances in the Baltic Sea
	PRF Cooperation Platform – Cooperation Platform on Port Reception Facilities in the Baltic Sea
	SAFE NAV – Group of Experts on Safety of Navigation
	SUBMERGED – Expert Group on Environmental Risks of Hazardous Submerged Objects
a.2 Region	nal Cooperation (RegionalCooperation)
The monitoring of this programme is:	

☑ Partly coordinated. Indicate missing component(s):☐ Coordinated monitoring is under development. Indicate by which group/project and by when a

recommendation on coordinated monitoring can be expected.

Common monitoring guidelines and quality assurance programme only partly in HELCOM COMBINE manual and also national. Common database is missing.

b. Monitoring strategies

b.1 Descriptor

☐ Fully coordinated

The programme supports the following obligatory MSFD Monitoring Strategies. Tick one or more relevant boxes.

⊠ D1	Biodiversity
□ D2	Non-indigenous Species
□ D3	Commercial fish and shellfish
□ D 4	Food webs
□ D 5	Eutrophication
⊠ D 6	Seafloor integrity
□ D7	Hydrographical conditions
□ D 8	Contaminants

□ D9	Contaminants in seafood			
□ D10	Marine litter			
□ D11	Energy including underwater noise			
b.2 BSAP se The sub-programn	egments ne serves the following BSAP segments. Tick one or more relevant boxes.			
⊠Eutrophication				
☐Hazardous sub	stances			
⊠Biodiversity				
☐Maritime activ	ities			
b.3 Monito	ring strategy description			
	etegy: Full description on monitoring of some of the relevant parameters, pobenthos and state of the substrate, available in HELCOM COMBINE manual. ation on monitoring frequency and spatial resolution from all countries is not			
b.4 BSAP Ed				
	cological objectives ost relevant option(s). Tick one or more boxes below.			
Choose only the m	ost relevant option(s). Tick one or more boxes below.			
Choose only the m	ost relevant option(s). Tick one or more boxes below. Concentrations of nutrients close to natural levels			
Choose only the m	□ Concentrations of nutrients close to natural levels □ Clear water			
Eutrophication	□ Concentrations of nutrients close to natural levels □ Clear water □ Natural level of algal blooms			
Eutrophication Hazardous	ost relevant option(s). Tick one or more boxes below. ☐ Concentrations of nutrients close to natural levels ☐ Clear water ☐ Natural level of algal blooms ☐ Natural distribution and occurrence of plants and animals			
Eutrophication	ost relevant option(s). Tick one or more boxes below. ☐ Concentrations of nutrients close to natural levels ☐ Clear water ☐ Natural level of algal blooms ☐ Natural distribution and occurrence of plants and animals ☐ Natural oxygen levels			
Eutrophication Hazardous	ost relevant option(s). Tick one or more boxes below. ☐ Concentrations of nutrients close to natural levels ☐ Clear water ☐ Natural level of algal blooms ☐ Natural distribution and occurrence of plants and animals ☐ Natural oxygen levels ☐ Concentrations of hazardous substances close to natural levels			
Eutrophication Hazardous	ost relevant option(s). Tick one or more boxes below. ☐ Concentrations of nutrients close to natural levels ☐ Clear water ☐ Natural level of algal blooms ☐ Natural distribution and occurrence of plants and animals ☐ Natural oxygen levels ☐ Concentrations of hazardous substances close to natural levels ☐ All fish safe to eat			
Eutrophication Hazardous	concentrations of nutrients close to natural levels Clear water Natural level of algal blooms Natural distribution and occurrence of plants and animals Natural oxygen levels Concentrations of hazardous substances close to natural levels All fish safe to eat Healthy wildlife			
Eutrophication Hazardous substances	cost relevant option(s). Tick one or more boxes below. Concentrations of nutrients close to natural levels Clear water Natural level of algal blooms Natural distribution and occurrence of plants and animals Natural oxygen levels Concentrations of hazardous substances close to natural levels All fish safe to eat Healthy wildlife Radioactivity at pre-Chernobyl levels			
Eutrophication Hazardous substances	concentrations of nutrients close to natural levels Clear water Natural level of algal blooms Natural distribution and occurrence of plants and animals Natural oxygen levels Concentrations of hazardous substances close to natural levels All fish safe to eat Healthy wildlife Radioactivity at pre-Chernobyl levels Natural landscapes and seascapes			
Eutrophication Hazardous substances Biodiversity Maritime	concentrations of nutrients close to natural levels Clear water Natural level of algal blooms Natural distribution and occurrence of plants and animals Natural oxygen levels Concentrations of hazardous substances close to natural levels All fish safe to eat Healthy wildlife Radioactivity at pre-Chernobyl levels Natural landscapes and seascapes Thriving and balanced communities of plants and animals			
Eutrophication Hazardous substances Biodiversity	concentrations of nutrients close to natural levels Clear water Natural level of algal blooms Natural distribution and occurrence of plants and animals Natural oxygen levels Concentrations of hazardous substances close to natural levels All fish safe to eat Healthy wildlife Radioactivity at pre-Chernobyl levels Natural landscapes and seascapes Thriving and balanced communities of plants and animals Viable populations of species			

\square No introductions of alien species from ships
\square Minimum air pollution from ships
☐ Zero discharges from offshore platforms
b.5 Gaps in monitoring In relation to the GES criteria addressed, indicate when sufficient monitoring was in place or by when sufficient coverage will be in place (Coverage_GEScriteria)
☐ Adequate monitoring was in place in 2014
☐ Adequate monitoring was in place by 2018
☐ Adequate monitoring is in place by July 2020
☐ Adequate monitoring will be in place by 2024
\square Monitoring is not being put in place for this descriptor due to a low risk
\square Monitoring for this descriptor is not relevant
Description of the implementation gaps and plans to complete the establishment and implementation of this descriptor monitoring strategy (Gaps_Plans):

There is very little monitoring data on the distribution of habitat-forming species currently available, whereas station or transect-based monitoring of the specific species is carried out in all the Baltic Sea countries. There are available modelling results for distribution of different habitat forming species Periodically mapping the bathymetry and distribution and condition of geological substrates is included in monitoring/inventory programmes in all the countries. No commonly agreed monitoring methods are in place, which target the extent and quality of the benthic habitats in the Baltic Sea. The drop-video technique in combination with traditional methods used for characterizing benthic communies (grab sampling, SCUBA diving) could be a promising and cost-effective solution at least for certain habitats. The need for joint and standardized monitoring methods/guidelines in the whole Baltic Sea area should be discussed and agreed, considering the diversity of natural conditions, environmental gradients as well as different reporting obligations.

c. Monitoring programmes

c.1 Purpose of monitoring

c.1a Assessment purpose in general

The programme supports the assessment of:

Tick the relevant box.

Temporal trends	Spatial distribution	State classification
		\boxtimes

The **programme** supports the assessment of: (MonitoringPurpose).

Note that the answer to this question will be decisive for whether to answer upcoming questions e.g.

upcoming questions on pressures should only be answered if the monitoring is defined as supporting the assessment of pressures.

Tick the relevant boxes.

Environmental state and impacts	Pressures in the marine environment	Pressures at source (land-based, riverine, sea-based ¹ and atmospheric sources)	Human activities causing the pressures	Effectiveness of measures
If this is selected fill in the following questions: c.1b	If this is selected fill in the following questions: c.1c, d	If this is selected fill in the following questions: c.1c, d		If this is selected fill in the following questions: c.1c, d

Give any other monitoring purpose e.g. if the programmes include supporting parameters for other monitoring programmes

Supporting HD assessments of marine habitats types.	

For questions 1b-1d, select when applicable for the sub-programme, the link from the Reporting on the 2020 update of Article 11 for the Marine Strategy Framework Directive (MSFD Guidance Document 17, 2020) (Features) to:

- Ecosystem components (relevant for monitoring and assessment for Article 8(1a) for D1C2-C5, D3, D4, D6C3-C5, D7C2)
- Pressures and impacts in the marine environment (relevant for monitoring and assessment for Article 8(1b) for D1C1, D2, D5, D6C1-C2, D7C1, D8, D9, D10, D11)
- Pressure inputs to the marine environment (relevant for monitoring and assessment for Article 10)
- Uses and human activities (relevant for monitoring and assessment for Article 8(1c) and 13)

c.1b • Ecosystem components (Features)

Choose only the most relevant option(s). Tick one or more boxes below.

Theme	Sub-theme	Label feature
Species	☐ Birds	☐ Grazing birds
		☐ Wading birds
		☐ Surface-feeding birds
		☐ Pelagic-feeding birds
		☐ Benthic-feeding birds
	☐ Mammals	☐ Small toothed cetaceans
		☐ Deep-diving toothed cetaceans

¹ Sea-based 'Pressures at source' refers to monitoring pressures from sea-based activities where the monitoring is directly at the activity rather than at a distance from or time period after it is generated by the activity (e.g. D1 incidental by-catch when fishing, D2 ballast water discharges, D6 use of bottom fishing gear, D8 contaminant discharges and pollution events from a vessel or pipeline, D11 impulsive sound events from a vessel or platform).

		☐ Baleen whales		
		☐ Seals		
	☐ Reptiles	☐ Turtles		
	☐ Fish	☐ Coastal fish		
		☐ Pelagic shelf fish		
		\square Demersal shelf fish		
		☐ Deep-sea fish		
		\square Commercially exploited fish and shellfish		
	\square Cephalopods	\square Coastal/shelf cephalopods		
		☐ Deep-sea cephalopods		
Habitats	☑ Benthic habitats	☑ Benthic broad habitats		
		☑ Other benthic habitats		
	☐ Pelagic habitats	☐ Pelagic broad habitats		
		☐ Other pelagic habitats		
Ecosystems	☐ Physical and hydrological	characteristics		
,				
	☐ Chemical characteristics			
	☐ Ecosystems, including	☐ Coastal ecosystems		
	food webs	☐ Shelf ecosystems		
		☐ Oceanic/deep-sea ecosystems		
	Pressures and impacts in to me most relevant option(s). Tick	the marine environment (Features) ck one or more boxes below.		
Theme	Label: Feature			
Biological	☐ Newly introduced non	-indigenous species		
	☐ Established non-indigenous species			
	☐ Species affected by inc	cidental by-catch		
Physical and				
hydrological	☑ Physical disturbance to	□ Physical disturbance to seabed		
	□ Physical loss of the seabed			
Substances,	☐ Eutrophication			
litter and	☐ Contaminants – non UPBT substances			
energy	☐ Contaminants − UPBT	☐ Contaminants − UPBT substances		
	☐ Contaminants – in sea	food		
	☐ Adverse effects on spe			
	<u></u>			

-		
_	☐ Acute pollution events	
_	☐ Litter in the environment	
-	☐ Impulsive sound in water	
	☐ Continuous low frequency sound	
c.1d • Pre	essure inputs to the marine environment (Features)	
Theme	Label: Feature	
Biological	☐ Input or spread of non-indigenous species	
	☐ Input of microbial pathogens	
	☐ Input of genetically modified species and translocation of native species	
_	\square Loss of, or change to, natural biological communities due to cultivation of animal or plant species	
_	$\hfill\Box$ Disturbance of species (e.g. where they breed, rest and feed) due to human presence	
	\Box Extraction of, or mortality/injury to, wild species (by commercial and recreational fishing and other activities)	
Substances,	☐ Input of nutrients — diffuse sources, point sources, atmospheric deposition	
litter and energy -	☐ Input of organic matter — diffuse sources and point sources	
chergy	\Box Input of other substances (e.g. synthetic substances, non-synthetic substances, radionuclides) — diffuse sources, point sources, atmospheric deposition, acute events	
	☐ Input of litter (solid waste matter, including micro-sized litter)	
	☐ Input of anthropogenic sound (impulsive, continuous)	
	$\hfill\Box$ Input of other forms of energy (including electromagnetic fields, light and heat)	
	☐ Input of water — point sources (e.g. brine)	
	es and human activities (Features) nost relevant option(s). Tick one or more boxes below.	
·		
Theme	Label: Feature	
Physical restructuring of	☐ Land claim	
rivers, coastline	☐ Canalisation and other watercourse modifications	
or seabed (water management)	☐ Coastal defence and flood protection	
	☐ Offshore structures (other than for oil/gas/renewables)	
	\square Restructuring of seabed morphology, including dredging and depositing of	

-	materials
Extraction of	☐ Extraction of minerals (rock, metal ores, gravel, sand, shell)
non-living resources	☐ Extraction of oil and gas, including infrastructure
	☐ Extraction of salt
	☐ Extraction of water
Production of energy	☐ Renewable energy generation (wind, wave and tidal power), including infrastructure
	□ Non-renewable energy generation
	☐ Transmission of electricity and communications (cables)
Extraction of	☐ Fish and shellfish harvesting (professional, recreational)
living resources	☐ Fish and shellfish processing
	☐ Marine plant harvesting
	☐ Hunting and collecting for other purposes
Cultivation of	☐ Aquaculture — marine, including infrastructure
living resources	☐ Aquaculture — freshwater
	☐ Agriculture
	☐ Forestry
Transport	☐ Transport infrastructure
	☐ Transport — shipping
	☐ Transport — air
	☐ Transport — land
Urban and	☐ Urban uses
industrial uses	☐ Industrial uses
	☐ Waste treatment and disposal
Tourism and	☐ Tourism and leisure infrastructure
leisure	☐ Tourism and leisure activities
Security/defence	☐ Military operations (subject to Article 2(2))
Education and research	☐ Research, survey and educational activities
c.2 Other leg The sub-programm one or more releva	e links with the following other international legislation (OtherPoliciesConventions). Tick
☐Bathing Water D	irective
☐Common Fisherie	es Policy and Data Collection Framework

⊠ Habitats Directive
☐ Birds Directive
□ Nitrates Directive
\square Urban Waste Water Treatment Directive
⊠Water Framework Directive
□ OSPAR Convention
☐Trilateral Wadden Sea Convention
□Other, Specify:
c.3 Implementation of Regional Cooperation (RegionalCooperation_implementation) Indicate the level of implementation by selecting one of the following:
⊠No coordination
☐ Agreed data collection methods
\square Common monitoring strategy (spatial and temporal design of programme)
\square Coordinated data collection (delivered separately by each country)
\Box Joint data collection (multinational delivery using same platform and/or algorithms)

c.4 Monitoring concepts

Monitoring concepts table²:

Current means of coordination	Features or elements Elements (Features) (Features_enum)	Parameter Parameters (Parameter) (ParametersOther)	Method MonitoringMethod (Monitoring Method) MonitoringMethodOt her)	QA/QC (Free text)	Frequency ³ MonitoringFrequency	Spatial resolution (density) of sampling (ProgrammeDescription)	Link to HELCOM core indicators ⁴ (RelatedIndicator) (RelatedIndicator_n ame	Spatial scope (SpatialSco pe)	Monitorin g started (year) (TemporalSc ope)	CPs monitoring ⁵ (CountryCode_E num)
National	Macroalgae (e.g. Fucus vesiculosus)	ABU/COV: Species abundance (numbers or coverage)	Drop-video mapping and verifying transects by divers.	National	Yearly, some areas once per 6 years	All WFD coastal water bodies		WFD CW	2014	EE
National	Angiosperms (Zostera and others)	DIST-R: Species distributional range/ pattern	Assessment of depth limits by video recording or diving along transects	National	Yearly (in rotation)	4 areas in each of the 5 sub- basins		Coastal	2021	FI
HELCOM	HabBenOthe r (Other benthic habitats) Soft-bottom macrofauna	ABU: Population size (abundance)	HEL-032 HELCOM COMBINE manual, Part C, Annex C8	HELCOM COMBINE manual	Yearly	3-10 offshore stations per subbasin and 3- 10 stations per coastal water type.		EEZ, coastal waters	1965	All HELCOM Contracting Parties

⁻

² Needed codelists can be found on 2020 update of Article 11 for the Marine Strategy Framework Directive (MSFD Guidance Document 17, 2020).

³ The option "Different for each country - see MORE overview" refers to the <u>overview</u> carried out in 2013

⁴ Give the name of HELCOM core indicators that are based on the monitoring parameter.

⁵ Provide information on the Contracting Partie(s) that are monitoring the parameter.

Current means of coordination	Features or elements	Parameter	Method	QA/QC	Frequency ³	Spatial resolution (density) of sampling	Link to HELCOM core indicators ⁴	Spatial scope	Monitorin g started (year)	CPs monitoring ⁵
National	Geological substrate	COV, EXT, DIST-P Coverage, extent and pattern						EEZ		All HELCOM Contracting Parties
National	Substrate condition	Abundance of drifting macroalgae on coarse and sandy substrates	drop video		yearly	Circa 20 sites (4 in 5 subbasins)		coastal waters	2021	FI

PARAMETER

Element/Parameter pair

Macroalgae / Species abundance

Angiosperms / Species distribution

Soft-bottom macrofauna / Population size

Macroalgae by drop videos / Macroalgae by drop videos

Angiosperms by drop videos / Angiosperms by drop videos

METHOD (Monitoring Details)

Macroalgae / Species abundance

In Estonia macroalagae species abundance is monitored according to WFD method (Torn&Martin, 2011).

Angiosperms / Species distribution

In Estonia species distribution of angiosperms is monitored according to WFD method (Torn&Martin, 2011).

Soft-bottom macrofauna / Population size

Drop video mapping, diving and the HELCOM COMBINE manual. Information on SE methods currently being used in the national MSFD Article 11 report (monitoring programme Vegetationsklädda bottnar): https://www.havochvatten.se/hav/samordning--

<u>fakta/miljoovervakning/marin-miljoovervakning/vegetationskladda-</u>bottnar.html

Macroalgae by drop videos / Macroalgae by drop videos

In Finland the monitoring has not started, however baseline mapping of the macroalgae distribution by drop video, diver transects and modeling has been carried out since 2004. The method for monitoring has been suggested.

In Denmark baseline mapping og reef sites (HD type 1170) and "bubbling reefs" (HD type: 1180) have been conducted in Natura 2000 sites including overall vegetation parameters (combined acoustic mapping with use of dropvideo, ROV and divers). No regular monitoring.

In Estonia drop camera observations are included in standard WFD monitoring procedure (Torn&Martin, 2011). Drop cameras are used also in monitoring of benthic habitats (Torn et al 2017).

In the German EEZ reefs are being monitored which includes macroalgae by drop camera and video transects and dredging

Angiosperms by drop videos / Angiosperms by drop videos

In Finland the monitoring will start in 2021 in all the 5 subbasins (4 areas in each) by yearly rotating cycle. In addition, baseline mapping of the angiosperm distribution by drop video, diver transects and modeling has been carried out since 2004.

In Denmark there are no regular monitoring but scientific projects working to test the applicability of drones, satelite images.

In Estonia drop camera observations are included in standard WFD monitoring procedure (Torn&Martin, 2011). Drop cameras are used also in monitoring of benthic habitats (Torn et al 2017).

In Germany there is a regular monitoring in inner and coastal waters

Substrate condition

In Finland substrate condition is evaluated from drop videos estimating abundance of loose-lying macroalgae on sandy and coarse substrates.

QA/QC

Macroalgae / Species abundance

National (EE)

Angiosperms / Species distribution

Soft-bottom macrofauna / Population size

National and HELCOM COMBINE manual

Macroalgae by drop videos / Macroalgae by drop videos

National

Angiosperms by drop videos / Angiosperms by drop videos

National

FREQUENCY

Frequency

Macroalgae / Species abundance

Estonia: yearly in 43 coastal waterbodies sites, at least once in 6 years in another 134 sites

Angiosperms / Species distribution

Estonia: yearly in 43 coastal waterbodies sites, at least once in 6 years in another 134 sites

Soft-bottom macrofauna / Population size

Estonia – once per year

Estonia: yearly in 4 areas, at least once in 6 years in another 12 areas; monitoring started in 1995.

Germany: yearly also video transects and dredging

Macroalgae by drop videos / Macroalgae by drop videos

Estonia: yearly in 4 coastal waterbodies, areas, at least once in 6 years in another 14 areas; use of drop videos started in 2005.

Germany: yearly also video transects and dredging

Angiosperms by drop videos / Angiosperms by drop videos

Finland: in ca 20 areas (4 in each 5 subbasins), monitored in yearly rotation.

Germany yearly (different methods)

Substrate condition

Finland: in ca 20 areas (4 in each 5 subbasins), monitored in yearly rotation.

SPATIAL SCOPE

Spatial Scope

Macroalgae / Species abundance and/or distribution

Estonia – transects in coastal WB-s

Angiosperms / Species distribution

Estonia – transects in coastal WB-s

Soft-bottom macrofauna / Population size

-

Macroalgae by drop videos / Macroalgae by drop videos

Mainly territorial waters but also in offshore areas (in German offshore waters Angiosperms do not occur)

Angiosperms by drop videos / Angiosperms by drop videos and other methods

Mainly territorial waters but also in offshore areas

Substrate condition

Finland: coastal waters

SPATIAL RESOLUTION (DENSITY) OF SAMPLING

Spatial resolution

Macroalgae / Species abundance

Angiosperms / Species distribution Soft-bottom macrofauna / Population size Estonia: spatial resolution - coastal waters WFD division - at least 3 transects in each surface water body Germany according to WFD in coastal waters and MSFD offshore + Habitas Directive on sandbanks and reefs Macroalgae by drop videos / Macroalgae by drop videos Estonia: spatial resolution - coastal waters WFD division - at least 3 transects in each surface water body Angiosperms by drop videos / Angiosperms by drop videos Finland: in ca 20 areas (4 in each 5 subbasins), several transects per area. Substrate condition

Finland: in ca 20 areas (4 in each 5 subbasins), several transects per site.

Provide considerations for the scale of aggregation of data for an indicator-based assessment Tick one or more relevant boxes below:

⊠HELCOM assessment unit Level 4: Subbasins with coastal WFD division
oxtimes HELCOM assessment unit Level 3: Subbasins with coastal and offshore division
\square HELCOM assessment unit Level 2: Subbasin
□HELCOM assessment unit Level 1: Baltic Sea
☐MSFD Region
□EU
\square Other (specify)
□Unknown

c.5 Monitoring and assessment requirements

Monitoring requirements:

The HELCOM core indicators directly linked to the sub-programme are still at a pre-core indicator stage of development. Creating specifications for monitoring requirements is a part of the development work of the indicators.

The monitoring of habitat-forming species and substrates focuses on covering wide sea areas with the purpose of providing data on extent and distribution. However, the data can also include parameters that enable state classification based on the condition of habitat-forming species. Monitoring of extent and distribution of habitatforming species and substrates, require spatial methods such as drop video, aerial surveys (aeroplanes, satellites, remote helicopters), multiple diving transects, randomized grab samples, multibeam sonars or different combinations of these methods. As the primary focus of monitoring is not in detecting temporal change but spatial status

(at certain time intervals), the monitoring frequency can be every 3 to 6 years.

The quality and quantity dimensions of habitat change are tradionally considered when assessing the status of habitats. There are hardly any operational methods used to estimate the quantity of a selected habitat that would also be useful for monitoring habitat extent (area or volume). To monitor habitat extent and describe the change of habitat extent or size statistically, methods that fully cover selected areas, delineate habitat boundaries, or use a large number of point observations may be used. There are several examples of methods that could qualify for this type of monitoring, including those applied on seagrass meadows being repeatedly mapped using aqua scope, video or remote sensing.

Although benthic monitoring for macrofauna and macrophytes is carried out within the COMBINE monitoring, no coordinated monitoring for habitat distribution and extent currently exists. Joint methods and guidelines will need to be agreed upon to provide data for the core indicators.

Adequacy for assessment of GES:

Monitoring should provide adequate data and information to enable the periodic assessment of environmental status, and distance from and progress towards GES as required by MSFD under Article 9 and Article 11.

	Yes	No
Adequate data?		
Established methods for assessment?		
Adequate understanding of GES?		\boxtimes
Adequate capacity to perform assessments?		

Assessment of natural variability

Qualitative. The borders of habitats and biotopes are naturally a continuum, which must be considered when assessing the dirstibution, extent and pattern of a biotope.

c.6 Data providers and access

From which database the data can be made available? Tick the relevant boxes below:

☐ HELCOM COMBINE	☐ HELCOM PLC	□HELCOM MORS
⊠Other:	National databa	ases

If the previous answer is "Other" please fill in the next questions (In case the answer is a HELCOM database, the HELCOM Secretariat will do it)

Data type Tick the releva	ant boxes below:				
□Unprocessed/raw Data					
⊠Processed Data sets					
☐ Data Products					
\square Modelled data					
Data management: General description of data management (DataManagement, Free text)					
<u>-</u>	will be used to make the data available? Tick the relevant boxes below and				
provide location (DataAcces					
☐ Providing URL to view d	ata:				
☐ Providing URL to downlo	oad data:				
☐ Provide location of data	in national data centre: Click here to enter text.				
☐ Provide location of data	in international data centre (e.g. RSC, ICES, EEA, EMODnet):				
	come available? (DataPublicationDate) , or even a past date if desired (MM/YYYY):				
EE: March next year after					
How frequently are the da	ta expected to be updated thereafter? Tick the relevant box below:				
□Every 6 years	□Weekly				
☐ Every 3 years	□Daily				
☐ Every 2 years	□Hourly				
⊠Yearly	□ Continually				
\Box 6-monthly	□One-off				
\square 3-monthly	☐ As needed				
\square Monthly	□Other (specify)				
\square 2-weekly	□Unknown				
List providing contact poin	ts in the Contracting Parties				
EE: Estonian Environmen	t Agency				
Has the data been used or	is it planned to be used in HELCOM assessments? Tick the relevant box below:				

⊠Yes	□No
Select if data is below:	s used in the following Baltic Sea Environment Fact Sheets (BSEF) Tick the relevant boxes
Biodiversity	
□Abundance	and distribution of marenzelleria species
☐ Abundance a	and distribution of Round goby
☐ Abundance a	and distribution of the Zebra mussel
\square Biopollution	level index
\square Observed no	on-indigenous and cryptogenic species in the Baltic Sea
☐ Population of	levelopment of Great Cormorant
☐ Population o	levelopment of Sandwich Tern
☐ Population of	levelopment of Southern Dunlin
□ Population □	Development of White-tailed Sea Eagle
☐Temporal de	velopment of Baltic coastal fish communities and key species
Eutrophicatio	on Control of the Con
\square Bacterioplan	nkton growth
\Box Chlorophyll-	a concentrations, temporal variations and regional differences from satellite remote sensing
\square Cyanobacter	ria biomass
\square Cyanobacter	rial blooms in the Baltic Sea
\square Cyanobacter	ria bloom index
\square Impacts of in	nvasive phytoplankton species on the Baltic Sea ecosystem in 1980-2008
□ Nitrogen atn	nospheric deposition to the Baltic Sea
□Nitrogen em	sissions to the air in the Baltic Sea area
□Phytoplankt	on biomass and species succession
\square Shifts in the	Baltic Sea summer phytoplankton communities in 1992-2006
☐Spatial distri	bution of the winter nutrient pool
☐Unusual phy	rtoplankton event
Hazardous su	bstances
□Atmospheric	deposition of heavy metals on the Baltic Sea
□Atmospheric	deposition of PCDD/Fs on the Baltic Sea
□Atmospheric	emissions of heavy metals in the Baltic Sea region
□Atmospherio	emissions of PCDD/Fs in the Baltic Sea region

\square Cesium-137 in B	altic Sea sediments			
\square Temporal trends in contaminants in Herring in the Baltic Sea in the period 1980-2010				
☐ Emissions from Baltic Sea shipping				
□ Illegal discharges	s of oil in the Baltic Sea			
\square Liquid discharge	s of Cs-137, Sr-90 and Co-60 into the Baltic Sea			
☐Trace metal cond	centrations and trends in Baltic surface and deep waters			
Hydrography				
\square Development of	Sea Surface Temperature in the Baltic Sea			
☐ Hydrography and	d Oxygen in the Deep Basins			
\square Ice season				
☐ Total and region	al runoff to the Baltic Sea			
\square Water Exchange	between the Baltic Sea and the North Sea, and conditions in the Deep Basins			
\square Wave climate in	the Baltic Sea			
	ost relevant option(s). Tick one or more boxes below.			
Descriptor 1	□ D1C1 – Primary:			
	The mortality rate per species from incidental by-catch is below levels which threaten the species, such that its long- term viability is ensured.			
	Member States shall establish the threshold values for the mortality rate from incidental by-catch per species, through regional or subregional cooperation.			
	☐ D1C2 — Primary:			
	The population abundance of the species is not adversely affected due to anthropogenic pressures, such that its long-term viability is ensured.			
	Member States shall establish threshold values for each species through regional or subregional cooperation, taking account of natural variation in population size and the mortality rates derived from D1C1, D8C4 and D10C4 and other relevant pressures. For species covered by Directive 92/43/EEC, these values shall be consistent with the Favourable Reference Population values established by the relevant Member States under Directive 92/43/EEC.			
	$\hfill\Box$ D1C3 — Primary for commercially- exploited fish and cephalopods and secondary for other species:			
	The population demographic characteristics (e.g. body size or age class structure, sex ratio, fecundity, and survival rates) of the species are indicative of a healthy population which is not adversely affected due to anthropogenic pressures.			
	Member States shall establish threshold values for specified characteristics of each			

species through regional or subregional cooperation, taking account of adverse effects

	on their health derived from D8C2, D8C4 and other relevant pressures.
	☐ D1C4 – Primary for species covered by Annexes II, IV or V to Directive 92/43/EEC and secondary for other species:
	The species distributional range and, where relevant, pattern is in line with prevailing physiographic, geographic and climatic conditions.
	Member States shall establish threshold values for each species through regional or subregional cooperation. For species covered by Directive 92/43/EEC, these shall be consistent with the Favourable Reference Range values established by the relevant Member States under Directive 92/43/EEC.
	☑ D1C5 – Primary for species covered by Annexes II, IV and V to Directive 92/43/EEC and secondary for other species:
	The habitat for the species has the necessary extent and condition to support the different stages in the life history of the species.
	□ D1C6 – Primary
	The condition of the habitat type, including its biotic and abiotic structure and its functions (e.g. its typical species composition and their relative abundance, absence of particularly sensitive or fragile species or species providing a key function, size structure of species), is not adversely affected due to anthropogenic pressures.
	Member States shall establish threshold values for the condition of each habitat type, ensuring compatibility with related values set under Descriptors 2, 5 and 8, through regional or subregional cooperation.
	1.08.01.01.01.08.01.01.00.01
Descriptor 2	□ D2C1 – Primary:
Descriptor 2	
Descriptor 2	□ D2C1 – Primary: The number of non-indigenous species which are newly introduced via human activity into the wild, per assessment period (6 years), measured from the reference year as reported for the initial assessment under Article 8(1) of Directive 2008/56/EC, is
Descriptor 2	□ D2C1 – Primary: The number of non-indigenous species which are newly introduced via human activity into the wild, per assessment period (6 years), measured from the reference year as reported for the initial assessment under Article 8(1) of Directive 2008/56/EC, is minimised and where possible reduced to zero. Member States shall establish the threshold value for the number of new introductions
Descriptor 2	□ D2C1 – Primary: The number of non-indigenous species which are newly introduced via human activity into the wild, per assessment period (6 years), measured from the reference year as reported for the initial assessment under Article 8(1) of Directive 2008/56/EC, is minimised and where possible reduced to zero. Member States shall establish the threshold value for the number of new introductions of non-indigenous species, through regional or subregional cooperation.
Descriptor 2	□ D2C1 − Primary: The number of non-indigenous species which are newly introduced via human activity into the wild, per assessment period (6 years), measured from the reference year as reported for the initial assessment under Article 8(1) of Directive 2008/56/EC, is minimised and where possible reduced to zero. Member States shall establish the threshold value for the number of new introductions of non-indigenous species, through regional or subregional cooperation. □ D2C2 — Secondary: Abundance and spatial distribution of established non-indigenous species, particularly of invasive species, contributing significantly to adverse effects on particular species
Descriptor 2	□ D2C1 − Primary: The number of non-indigenous species which are newly introduced via human activity into the wild, per assessment period (6 years), measured from the reference year as reported for the initial assessment under Article 8(1) of Directive 2008/56/EC, is minimised and where possible reduced to zero. Member States shall establish the threshold value for the number of new introductions of non-indigenous species, through regional or subregional cooperation. □ D2C2 — Secondary: Abundance and spatial distribution of established non-indigenous species, particularly of invasive species, contributing significantly to adverse effects on particular species groups or broad habitat types.
Descriptor 2	□ D2C1 − Primary: The number of non-indigenous species which are newly introduced via human activity into the wild, per assessment period (6 years), measured from the reference year as reported for the initial assessment under Article 8(1) of Directive 2008/56/EC, is minimised and where possible reduced to zero. Member States shall establish the threshold value for the number of new introductions of non-indigenous species, through regional or subregional cooperation. □ D2C2 — Secondary: Abundance and spatial distribution of established non-indigenous species, particularly of invasive species, contributing significantly to adverse effects on particular species groups or broad habitat types. □ D2C3 — Secondary: Proportion of the species group or spatial extent of the broad habitat type which is adversely altered due to non-indigenous species, particularly invasive non-indigenous
Descriptor 2 Descriptor 3	□ D2C1 − Primary: The number of non-indigenous species which are newly introduced via human activity into the wild, per assessment period (6 years), measured from the reference year as reported for the initial asessment under Article 8(1) of Directive 2008/56/EC, is minimised and where possible reduced to zero. Member States shall establish the threshold value for the number of new introductions of non-indigenous species, through regional or subregional cooperation. □ D2C2 — Secondary: Abundance and spatial distribution of established non-indigenous species, particularly of invasive species, contributing significantly to adverse effects on particular species groups or broad habitat types. □ D2C3 — Secondary: Proportion of the species group or spatial extent of the broad habitat type which is adversely altered due to non-indigenous species, particularly invasive non-indigenous species. Member States shall establish the threshold values for the adverse alteration to species groups and broad habitat types due to non-indigenous species, through regional or

	scientific bodies shall be consulted in accordance with Article 26 of Regulation (EU) No 1380/2013.
	□ D3C2 — Primary:
	The Spawning Stock Biomass of populations of commercially-exploited species are above biomass levels capable of producing maximum sustainable yield. Appropriate scientific bodies shall be consulted in accordance with Article 26 of Regulation (EU) No 1380/2013.
	□ D3C3 — Primary:
	The age and size distribution of individuals in the populations of commercially-exploited species is indicative of a healthy population. This shall include a high proportion of old/large individuals and limited adverse effects of exploitation on genetic diversity.
	Member States shall establish threshold values through regional or subregional cooperation for each population of species in accordance with scientific advice obtained pursuant to Article 26 of Regulation (EU) No 1380/2013.
Descriptor 4	□ D4C1 — Primary:
	The diversity (species composition and their relative abundance) of the trophic guild is not adversely affected due to anthropogenic pressures.
	Member States shall establish threshold values through regional or subregional cooperation.
	□ D4C2 — Primary:
	The balance of total abundance between the trophic guilds is not adversely affected due to anthropogenic pressures.
	Member States shall establish threshold values through regional or subregional cooperation.
	□ D4C3 — Secondary:
	The size distribution of individuals across the trophic guild is not adversely affected due to anthropogenic pressures.
	Member States shall establish threshold values through regional or subregional cooperation.
	\square D4C3 — Secondary (to be used in support of criterion D4C2, where necessary):
	Productivity of the trophic guild is not adversely affected due to anthropogenic pressures.
	Member States shall establish threshold values through regional or subregional cooperation.
Descriptor 5	□ D5C1 — Primary:
	Nutrient concentrations are not at levels that indicate adverse eutrophication effects.
	The threshold values are as follows:
	(a) in coastal waters, the values set in accordance with Directive 2000/60/EC;
	(b) beyond coastal waters, values consistent with those for coastal waters under Directive 2000/60/EC. Member States shall establish those values through

regional or subregional cooperation
□ D5C2 — Primary:
Chlorophyll a concentrations are not at levels that indicate adverse effects of nutrient enrichment.
The threshold values are as follows:
(c) in coastal waters, the values set in accordance with Directive 2000/60/EC;
(d) beyond coastal waters, values consistent with those for coastal waters under Directive 2000/60/EC. Member States shall establish those values through regional or subregional cooperation.
☐ D5C3 — Secondary:
The number, spatial extent and duration of harmful algal bloom events are not at levels that indicate adverse effects of nutrient enrichment.
☐ D5C4 — Secondary:
The photic limit (transparency) of the water column is not reduced, due to increases in suspended algae, to a level that indicates adverse effects of nutrient enrichment.
The threshold values are as follows:
(e) in coastal waters, the values set in accordance with Directive 2000/60/EC;
(f) beyond coastal waters, values consistent with those for coastal waters under Directive 2000/60/EC. Member States shall establish those values through regional or subregional cooperation.
\square D5C5 — Primary (may be substituted by D5C8):
The concentration of dissolved oxygen is not reduced, due to nutrient enrichment, to levels that indicate adverse effects on benthic habitats (including on associated biota and mobile species) or other eutrophication effects.
The threshold values are as follows:
(g) in coastal waters, the values set in accordance with Directive 2000/60/EC;
(h) beyond coastal waters, values consistent with those for coastal waters under Directive 2000/60/EC. Member States shall establish those values through regional or subregional cooperation.
□ D5C6 — Secondary:
The abundance of opportunistic macroalgae is not at levels that indicate adverse effects of nutrient enrichment.
The threshold values are as follows:
(a) in coastal waters, the values set in accordance with Directive 2000/60/EC;
(b) should this criterion be relevant for waters beyond coastal waters, values consistent with those for coastal waters under Directive 2000/60/EC. Member States shall establish those values through regional or subregional cooperation.
□ D5C7 — Secondary:
The species composition and relative abundance or depth distribution of macrophyte

communities achieve values that indicate there is no adverse effect due to nutrient enrichment including via a decrease in water transparency, as follows: (a) in coastal waters, the values set in accordance with Directive 2000/60/EC; (b) should this criterion be relevant for waters beyond coastal waters, values consistent with those for coastal waters under Directive 2000/60/EC. Member States shall establish those values through regional or subregional cooperation. ☐ D5C8 — Secondary: (except when used as a substitute for D5C5): The species composition and relative abundance of macrofaunal communities, achieve values that indicate that there is no adverse effect due to nutrient and organic enrichment, as follows: (a) in coastal waters, the values for benthic biological quality elements set in accordance with Directive 2000/60/EC; (b) beyond coastal waters, values consistent with those for coastal waters under Directive 2000/60/EC. Member States shall establish those values through regional or subregional cooperation. Descriptor 6 \square D6C1 – Primary: Spatial extent and distribution of physical loss (permanent change) of the natural seabed. \square D6C2 – Primary: Spatial extent and distribution of physical disturbance pressures on the seabed. \boxtimes D6C3 – Primary: Spatial extent of each habitat type which is adversely affected, through change in its biotic and abiotic structure and its functions (e.g. through changes in species composition and their relative abundance, absence of particularly sensitive or fragile species or species providing a key function, size structure of species), by physical disturbance. Member States shall establish threshold values for the adverse effects of physical disturbance, through regional or subregional cooperation. \boxtimes D6C4 – Primary: The extent of loss of the habitat type, resulting from anthropogenic pressures, does not exceed a specified proportion of the natural extent of the habitat type in the assessment area. Member States shall establish the maximum allowable extent of habitat loss as a proportion of the total natural extent of the habitat type, through cooperation at Union level, taking into account regional or subregional specificities. ☑ D6C5 – Primary: The extent of adverse effects from anthropogenic pressures on the condition of the habitat type, including alteration to its biotic and abiotic structure and its functions (e.g. its typical species composition and their relative abundance, absence of particularly sensitive or fragile species or species providing a key function, size structure of species), does not exceed a specified proportion of the natural extent of the habitat type in the assessment area.

	Member States shall establish threshold values for adverse effects on the condition of each habitat type, ensuring compatibility with related values set under Descriptors 2, 5, 6, 7 and 8, through cooperation at Union level, taking into account regional or subregional specificities. Member States shall establish the maximum allowable extent of those adverse effects as a proportion of the total natural extent of the habitat type, through cooperation at Union level, taking into account regional or subregional specificities.
Descriptor 7	□ D7C1 – Secondary:
	Spatial extent and distribution of permanent alteration of hydrographical conditions (e.g. changes in wave action, currents, salinity, temperature) to the seabed and water column, associated in particular with physical loss(1) of the natural seabed.
	□ D7C2 – Secondary:
	Spatial extent of each benthic habitat type adversely affected (physical and hydrographical characteristics and associated biological communities) due to permanent alteration of hydrographical conditions.
Descriptor 8	□ D8C1 – Primary:
	Within coastal and territorial waters, the concentrations of contaminants do not exceed the following threshold values:
	(a) for contaminants set out under point 1(a) of criteria elements, the values set in accordance with Directive 2000/60/EC;
	(b) when contaminants under point (a) are measured in a matrix for which no value is set under Directive 2000/60/EC, the concentration of those contaminants in that matrix established by Member States through regional or subregional cooperation;
	(c) for additional contaminants selected under point 1(b) of criteria elements, the concentrations for a specified matrix (water, sediment or biota) which may give rise to pollution effects. Member States shall establish these concentrations through regional or subregional cooperation, considering their application within and beyond coastal and territorial waters.
	Beyond territorial waters, the concentrations of contaminants do not exceed the following threshold values:
	(a) for contaminants selected under point 2(a) of criteria elements, the values as applicable within coastal and territorial waters;
	(b) for contaminants selected under point 2(b) of criteria elements, the concentrations for a specified matrix (water, sediment or biota) which may give rise to pollution effects. Member States shall establish these concentrations through regional or subregional cooperation.
	□ D8C2 – Secondary:
	The health of species and the condition of habitats (such as their species composition and relative abundance at locations of chronic pollution) are not adversely affected due to contaminants including cumulative and synergetic effects.
	Member States shall establish those adverse effects and their threshold values through

	regional or subregional cooperation.
	□ D8C3 – Primary:
	The spatial extent and duration of significant acute pollution events are minimised.
	\square D8C4 – Secondary (to be used when a significant acute pollution event has occurred):
	The adverse effects of significant acute pollution events on the health of species and on the condition of habitats (such as their species composition and relative abundance) are minimised and, where possible, eliminated.
Descriptor 9	□ D9C1 – Primary:
	The level of contaminants in edible tissues (muscle, liver, roe, flesh or other soft parts, as appropriate) of seafood (including fish, crustaceans, molluscs, echinoderms, seaweed and other marine plants) caught or harvested in the wild (excluding fin-fish from mariculture) does not exceed:
	 (a) for contaminants listed in Regulation (EC) No 1881/2006, the maximum levels laid down in that Regulation, which are the threshold values for the purposes of this Decision;
	(b) for additional contaminants, not listed in Regulation (EC) No 1881/2006, threshold values, which Member States shall establish through regional or subregional cooperation.
Descriptor 10	□ D10C1 – Primary:
	The composition, amount and spatial distribution of litter on the coastline, in the surface layer of the water column, and on the seabed, are at levels that do not cause harm to the coastal and marine environment.
	Member States shall establish threshold values for these levels through cooperation at Union level, taking into account regional or subregional specificities.
	□ D10C2 — Primary:
	The composition, amount and spatial distribution of micro-litter on the coastline, in the surface layer of the water column, and in seabed sediment, are at levels that do not cause harm to the coastal and marine environment.
	Member States shall establish threshold values for these levels through cooperation at Union level, taking into account regional or subregional specificities.
	□ D10C3 — Secondary:
	The amount of litter and micro-litter ingested by marine animals is at a level that does not adversely affect the health of the species concerned. Member States shall establish threshold values for these levels through regional or subregional cooperation.
	□ D10C4 — Secondary:
	The number of individuals of each species which are adversely affected due to litter, such as by entanglement, other types of injury or mortality, or health effects. Member States shall establish threshold values for the adverse effects of litter, through regional or subregional cooperation.

Descriptor 11	□ D11C1 – Primary:
	The spatial distribution, temporal extent, and levels of anthropogenic impulsive sound sources do not exceed levels that adversely affect populations of marine animals.
	Member States shall establish threshold values for these levels through cooperation at Union level, taking into account regional or subregional specificities.
	□ D11C2 – Primary:
	The spatial distribution, temporal extent and levels of anthropogenic continuous low-frequency sound do not exceed levels that adversely affect populations of marine animals.
	Member States shall establish threshold values for these levels through cooperation at Union level, taking into account regional or subregional specificities.

d. References

Make a list of cited references and literature for further supportive information.

HELCOM (2013) HELCOM Underwater Biotope and habitat classification system (HELCOM HUB)

Torn, Kaire; Martin, Georg (2011). Assessment method for the ecological status of Estonian coastal waters based on submerged aquatic vegetation. In: Brebbia, C.A.; Beriatos, E. (Ed.). Sustainable Development and Planning V (443–452).. Southampton: WIT Press. (WIT Transactions on Ecology and the Environment; 150).10.2495/SDP110371.

Torn, K.; Herkül, K.; Martin, G.; Oganjan, K. (2017). Assessment of quality of three marine benthic habitat types in northern Baltic Sea. Ecological Indicators, 73, 772–783.10.1016/j.ecolind.2016.10.037.

Germany:

https://www.bfn.de/fileadmin/MDB/documents/themen/meeresundkuestenschutz/downloads/Monitoringberichte/BfN-Monitoring-Benthos-Ostsee-2009.pdf